SPUTUM

Synonyms

- Consecutive Early Morning Sputum
- Post Bronchoscopy Sputum

Test Code: 9012 / **CPT Code:** 88112

Supplies:

- 1. Non-sterile specimen containers with 30ml of CytoLyt
- 2. Yellow Non-Gyn requisitions
- 3. Biohazard Bags

Collection:

- 1. USE UNIVERSAL PRECAUTIONS
- 2. For consecutive early morning sputum samples:
 - Collect early morning, "deep cough" specimen (not saliva) and place into CytoLyt container.
 - Specimen should be obtained before the patient has eaten.
- 3. For post bronchoscopy sputum samples:
 - Collect separate sputum specimens at 24 and 48 hours after bronchoscopy is performed.
 - Collect "deep cough" specimen (not saliva) and place into CytoLyt container.
- 4. Secure the lid and gently shake specimen.
- 5. It is important to keep each specimen separate and state the collection date on both the specimen and the requisition card.
- 6. Secure the lid and label specimen container with the following:
 - Patient's name
 - Physician's name
 - Source of specimen
 - Date of collection
- 7. Complete the yellow Non-Gyn requisition card with the following information:
 - Patient's name, birth date and sex
 - Physician's name and clinic account number
 - Source of specimen
 - Date of collection
 - All pertinent clinical information (especially history of cancer, chemotherapy and radiation treatments).
 - A separate requisition card must be filled out for each specimen.
- 8. Send specimen container and requisition card in a Biohazard bag to DCCC.
 - Tighten lid of CytoLyt container, place the specimen in the ziplock portion of the bag and seal it.
 - Place the yellow Non-Gyn requisition card into the back pocket with the overlapping flaps.

Storage Instructions:

1. If more than a 24-hr delay is anticipated between collection and receipt in the laboratory, please refrigerate specimen.